

Provided for non-commercial research and educational use. Not for reproduction, distribution or commercial use.
--

Serdica

Mathematical Journal

Сердика

Математическо списание

The attached copy is furnished for non-commercial research and education use only.
Authors are permitted to post this version of the article to their personal websites or institutional repositories and to share with other researchers in the form of electronic reprints.
Other uses, including reproduction and distribution, or selling or licensing copies, or posting to third party websites are prohibited.

For further information on
Serdica Mathematical Journal
which is the new series of
Serdica Bulgaricae Mathematicae Publicationes
visit the website of the journal <http://www.math.bas.bg/~serdica>
or contact: Editorial Office
Serdica Mathematical Journal
Institute of Mathematics and Informatics
Bulgarian Academy of Sciences
Telephone: (+359-2)9792818, FAX:(+359-2)971-36-49
e-mail: serdica@math.bas.bg

EQUICONVERGENCE AND EQUISUMMABILITY OF JACOBI SERIES

Georgi Boychev

Communicated by O. Mushkarov

ABSTRACT. In this paper we give some results concerning the equiconvergence and equisummability of series in Jacobi polynomials.

1. Jacobi polynomials and series in the complex plane. Let $\alpha > -1$ and $\beta > -1$. The polynomials $\left\{P_n^{(\alpha,\beta)}(z)\right\}_{n=0}^{+\infty}$ defined by the equalities

$$P_n^{(\alpha,\beta)}(z) = \binom{n+\alpha}{n} F\left(-n, n+\alpha+\beta+1, \alpha+1; \frac{1-z}{2}\right),$$

$$z \in \mathbb{C}, \quad n = 0, 1, 2, \dots,$$

where $F(a, b, c; \zeta)$ is the Gauss hypergeometric function, are called *Jacobi polynomials with parameters α and β* .

2010 *Mathematics Subject Classification*: 33C45, 40G05.

Key words: Jacobi polynomials, Jacobi series, asymptotic formula, convergence, summability, equisummability, equiconvergence.

There exists a unique complex function h , holomorphic in the region $G = \mathbb{C} \setminus [-1, +1]$, such that $h^2(z) = z^2 - 1$ when $z \in G$ and $h(x) > 0$ when $x > 1$. The value of this function at any point $z \in G$ is denoted by $\sqrt{z^2 - 1}$. The function w defined in G as

$$w(z) = z + \sqrt{z^2 - 1}$$

is holomorphic in G . Moreover, $w(z) \neq 0$ and $[w(z) + (w(z))^{-1}]/2 = z$ when $z \in G$. Therefore, $w(z)$ may be considered as an inverse of the Zhukovsky function. As it is well known, the latter one is univalent in the region $D = \{w : |w(z)| > 1\}$ and maps it onto G . Hence, the function w maps G onto D . The function w is meromorphic in the region $G = \overline{\mathbb{C}} \setminus [-1, +1]$ and $\lim_{z \rightarrow \infty} w(z) = \infty$.

G. Szegő gives an asymptotic expansion for Jacobi polynomials in the region G [4, Theorem 8.21.1]. From this expansion it follows the asymptotic formula ($n \geq 1$, $p = 0, 1, 2, \dots$)

$$(1.1) \quad P_n^{(\alpha, \beta)}(z) = \psi(z)n^{-1/2}(w(z))^n \left\{ \sum_{k=0}^p \varphi_k(z)n^{-k} + \varphi_{n,p}(z) \right\},$$

where $\psi(z) \neq 0$, $\{\varphi_k(z)\}_{k=0}^{+\infty}$, $\{\varphi_{n,p}(z)\}_{n=1}^{+\infty}$ ($p = 0, 1, 2, \dots$) are a complex functions holomorphic in the region G and

$$(1.2) \quad \varphi_{n,p}(z) = O(n^{-p-1}) \quad (n \rightarrow \infty)$$

uniformly on every compact subset of G . In particular, $\varphi_0(z) \equiv 1$.

If $p = 0$, then it follows from (1.1) and (1.2) that [3, (II.3.1)]

$$(1.3) \quad P_n^{(\alpha, \beta)}(z) = \psi(z)n^{-1/2}(w(z))^n \{1 + \varphi_{n,0}(z)\},$$

where $\varphi_{n,0}(z) = O(n^{-1})$ ($n \rightarrow \infty$) uniformly on every compact subset of G .

Suppose that $r > 1$. Denote by $\gamma(r)$ the image of the circle $C(0, r) = \{z \in \mathbb{C} : z = r \exp i\theta, 0 \leq \theta \leq 2\pi\}$ by the Zhukovsky transformation. As it is well known, $\gamma(r)$ is the ellipse with focuses at the points -1 and $+1$ and semiaxes $(r + r^{-1})/2$ and $(r - r^{-1})/2$.

Let $\overline{E(r)}$ be the interior of $\gamma(r)$. Further, we denote $E(\infty) = \mathbb{C}$, $E(1) = \emptyset$, $E^*(r) = \overline{\mathbb{C}} \setminus \overline{E(r)}$ ($1 < r < \infty$), $E^*(\infty) = \emptyset$ and $E^*(1) = G$.

The series of the kind

$$(1.4) \quad \sum_{n=0}^{+\infty} a_n P_n^{(\alpha, \beta)}(z)$$

has the name *Jacobi series*.

Theorem 1.1 [3, (IV.1.1), (a)]. *If the series (1.4) converges at a point $z_0 \in G$, then it is absolutely uniformly convergent on every compact subset of the region $E(r)$ with $r = |w(z_0)|$.*

We remark that in the proof of this theorem it is used the asymptotic formula (1.3).

Theorem 1.2 [3, (IV.1.1), (b)]. *If*

$$(1.5) \quad r = \max \left\{ 1, \left(\lim_{n \rightarrow +\infty} \sup |a_n|^{1/n} \right)^{-1} \right\},$$

then the series (1.4) is absolutely uniformly convergent on every compact subset of the domain $E(r)$ and diverges in $E^(r)$.*

The equality (1.5) can be regarded as a *formula of Cauchy-Hadamard type* for the Jacobi series. If $r > 1$, then the sum of the series (1.4) is a complex function, holomorphic in the domain $E(r)$.

2. (C, δ) -summability of a series. Let us remind that a series

$$(2.1) \quad \sum_{n=0}^{+\infty} u_n$$

is said to be (C, δ) -summable for $\delta > -1$, if there exists

$$\sigma = \lim_{n \rightarrow +\infty} (A_n^\delta)^{-1} \sum_{k=0}^n A_{n-k}^{\delta-1} s_k,$$

where $A_k^\delta = \Gamma(k + \delta + 1) / [\Gamma(k + 1)\Gamma(\delta + 1)]$ ($\delta > -1$; $k = 0, 1, \dots$) and $\{s_k\}_{k=0}^{+\infty}$ are the partial sums of the series (2.1).

Usually the complex number σ is called (C, δ) -sum of the series (2.1). Every convergent series is (C, δ) -summable for every $\delta > 0$ and its (C, δ) -sum is equal to its sum in the usual sense.

3. Equisummability of Jacobi series. Let us to consider a series of kind

$$(3.1) \quad \sum_{n=1}^{+\infty} a_n n^{-1/2} (w(z))^n.$$

In the following proposition we give a relation between the series (1.4) and (2.1).

Theorem 3.1. *Let $\delta > -1$ and $z_0 \in G$. Then the series (1.4) is (C, δ) -summable at z_0 if and only if the series (2.1) is (C, δ) -summable at z_0 .*

Proof. Let p be an integer such that $p > \delta + 1$. Further, we assume that $a_0 = 0$.

Suppose that the series (1.4) is (C, δ) -summable for $z = z_0$. Applying [2, Theorem 46], we obtain that

$$(3.2) \quad a_n P_n^{(\alpha, \beta)}(z_0) = o(n^\delta) \quad (n \rightarrow \infty).$$

We define the numbers $\mu_1, \mu_2, \dots, \mu_p$ by the following equalities

$$\mu_1 = \varphi_1(z_0)$$

$$\mu_2 = \varphi_2(z_0) - \mu_1 \varphi_1(z_0)$$

.....

$$\mu_p = \varphi_p(z_0) - \sum_{j=1}^{p-1} \varphi_{p-j}(z_0) \mu_j.$$

If

$$b_n = a_n P_n^{(\alpha, \beta)}(z_0) \{1 - \mu_1 n^{-1} - \dots - \mu_p n^{-p}\},$$

then from [2, Theorem 74] it follows that the series $\sum_{k=1}^{+\infty} b_n$ is (C, δ) -summable.

Using the asymptotic formulas (1.1) and (1.2), we conclude that

$$b_n = \psi(z_0) a_n n^{-1/2} (w(z_0))^n \{1 + d_{n,p}(z_0)\},$$

where $d_{n,p}(z_0) = O(n^{-p-1})$ ($n \rightarrow \infty$). Obviously, $\psi(z_0) \neq 0$ and $P_n^{(\alpha, \beta)}(z_0) \neq 0$ ($n = 1, 2, \dots$). Then, from (3.2), (1.1) and (1.2) it follows that

$$a_n \psi(z_0) n^{-1/2} (w(z_0))^n = o(n^\delta) \quad (n \rightarrow \infty).$$

Further, we have

$$q_n = \psi(z_0) a_n n^{-1/2} (w(z_0))^n d_{n,p}(z_0) = o(n^\delta) O(n^{-p-1}) \quad (n \rightarrow \infty).$$

Since $p > \delta + 1$ then

$$q_n = O(n^{-2}) \quad (n \rightarrow \infty).$$

Hence, the series $\sum_{k=1}^{+\infty} q_n$ is convergent. Applying [3, Theorem 45] we deduce that this series is (C, δ) -summable. Then, from the equalities

$$a_n n^{-1/2} (w(z_0))^n = b_n / \psi(z_0) - q_n \quad (n = 1, 2, \dots)$$

it follows that the series (3.1) is (C, δ) -summable.

Now, let us assume that the series (3.1) is (C, δ) -summable for $z = z_0$. Then

$$a_n n^{-1/2} (w(z_0))^n = o(n^\delta) \quad (n \rightarrow \infty).$$

This asymptotic formula and the asymptotic formula (1.2) yield

$$c_n = \psi(z_0) a_n n^{-1/2} (w(z_0))^n d_{n,p}(z_0) = o(n^\delta) O(n^{-p-1}).$$

Since $p > \delta + 1$, then $c_n = O(n^{-2})$. Hence the series $\sum_{k=1}^{+\infty} c_n$ converges. According to [3, Theorem 45] it is (C, δ) -summable.

The series of the kind

$$\sum_{n=1}^{+\infty} a_n (w(z_0))^n n^{-s-1/2},$$

where $s = 1, 2, \dots, p$, is (C, δ) -summable by Theorem 45 from [3]. Then, from the equalities $(n = 1, 2, \dots)$

$$a_n P_n^{(\alpha, \beta)}(z_0) = \psi(z_0) a_n n^{-1/2} (w(z_0))^n + \psi(z_0) \sum_{s=1}^p a_n (w(z_0))^n \varphi_s(z_0) n^{-s-1/2} + c_n$$

it follows that the series (3.1) is (C, δ) -summable for $z = z_0$. Thus Theorem 3.1 is proved. \square

Having in mind the proof of Theorem 3.1 it is easy to establish the following result:

Theorem 3.2. *If $\delta > -1$ and K is a compact subset of the region G , then the series (1.4) is uniformly (C, δ) -summable on K if and only if the series (3.1) is uniformly (C, δ) -summable on K .*

The following statement is a corollary of Theorem 3.1.

Theorem 3.3. *If $z_0 \in G$, then the series (1.4) converges for $z = z_0$ if and only if the series (3.1) converges for $z = z_0$.*

Theorems 3.1 and 3.3 can be called *Theorems for equisummability and equiconvergence* of Jacobi series, respectively.

4. Fatou-Riesz theorem for Jacobi series. Let

$$(4.1) \quad \sum_{n=0}^{+\infty} a_n t^n$$

be a power series with finite and different from zero radius r of convergence and $g(t)$ be a sum of (4.1) in the region $\text{int}\{C(0, r)\}$. It is well known that if the coefficients $\{a_n\}_{n=0}^{+\infty}$ satisfy certain additional conditions, there is a relation between the regular points of the function $g(t)$ on the circumference $C(0, r)$ and the behaviour of (4.1) on the same circumference. The following result holds [6, (9.21)].

Theorem 4.1 (Fatou-Riesz). *If the power series (4.1) has a finite radius r of convergence and $a_n r^n = o(n^\delta)$ ($n \rightarrow +\infty$) for some $\delta > -1$, then the series (4.1) is uniformly (C, δ) -summable on every close arc of $C(0, r)$, all points of which are regular for the function $f(t)$.*

V. Yatsun in [5] described the relations between the singular points of Jacobi series (1.5) and those of the corresponding power series (4.1). In particular it follows from his main statement:

Theorem 4.2. *Let $1 < r = \lim_{n \rightarrow +\infty} \sup |a_n|^{1/n} < +\infty$ and $z_0 \in \gamma(r)$ be a regular point for the sum $f(z)$ of (1.4) in $E(r)$. Then $t_0 = w(z_0) \in C(0, r)$ is a regular point for the sum $g(t)$ of (4.1).*

Using Theorems 4.1 and 4.2 we are going to prove a theorem for the Jacobi series, which is similar to Theorem 4.1.

Theorem 4.3 [1]. *Suppose that r from (1.4) satisfies the inequalities $1 < r < +\infty$ and $a_n r^n = o(n^\delta)$ ($n \rightarrow +\infty$) for some $\delta > -1/2$. Then the series*

(1.4) is uniformly $(C, \delta - 1/2)$ -summable on every close arc $\gamma_1 \subset \gamma(r)$, all points of which are regular for the sum $f(z)$ of (1.4).

Proof. Let $\gamma_1^* = \{t : t = w(z), z \in \gamma_1\}$. Obviously, γ_1^* is a close arc of the circumference $C(0, r)$. According to Theorem 4.2 all point of this arc are regular for the sum $g(t)$ of (4.1). Then, using Theorem 4.1 the series (4.1), we obtain that this series is uniformly (C, δ) -summable on the arc γ_1^* . Then, the series

$$\sum_{n=1}^{+\infty} a_n t^n$$

is also uniformly (C, δ) -summable on the arc γ_1^* . Using [2, Theorem 76], we obtain that the series

$$\sum_{n=1}^{+\infty} a_n n^{-1/2} t^n$$

is uniformly $(C, \delta - 1/2)$ -summable on the arc γ_1^* . Hence, the series (3.1) is uniformly $(C, \delta - 1/2)$ -summable for $z \in \gamma_1$. According Theorem 3.2 the series (1.5) is uniformly $(C, \delta - 1/2)$ -summable on the arc γ_1 . Thus Theorem 4.3 is proved. \square

Remark. The proof of Theorem 4.3 is considerably shorter than its proof given in [1].

REFERENCES

- [1] G. BOYCHEV. Fatou-Riesz's theorem for series in Jacobi polynomials. *C. R. Acad. Bulgare Sci.* **37**, 4 (1984), 427–428.
- [2] G. HARDY. Divergent Series. Oxford, At the Clarendon Press (Geoffrey Cumberlege), 1949.
- [3] P. RUSEV. Classical Orthogonal Polynomials and Their Associated Functions in Complex Plane. Prof. Marin Drinov Academic Publshing House, Sofia, 2005.
- [4] G. SZEGÖ. Orthogonal Polynomials. AMS Colloquium Publ. vol. **23**, New York, American Mathematical Society, 1939.

- [5] V. YATSUN. Analytic continuation of expansions in Jacobi polynomials. *Ukrain. Math. Zh.* **21**, 4 (1969), 511–521 (in Russian); English translation in *Ukr. Math. J.* **21** (1969), 436–444.
- [6] A. ZYGMUND. Trigonometric series. Volumes I and II, 3rd ed. Cambridge Mathematical Library. Cambridge, Cambridge University Press, 2002.

Trakia University
6000 Stara Zagora, Bulgaria
e-mail: GBoychev@hotmail.com

Received February 24, 2012